

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

35-1073: Polyclonal Antibody to HER2 (Phospho-Tyr877)

Clonality: Polyclonal Application: WB,IHC,IF

Reactivity: Human, Mouse, Rat

Gene : ERBB2
Gene ID : 2064
Uniprot ID : P04626
Format : Purified

Alternative Name : C-erbB-2, ErbB2 Isotype : Rabbit IgG

Immunogen Information: Peptide sequence around phosphorylation site of tyrosine 877 (T-E-Y(p)-H-A) derived from Human

HER2.

Description

Essential component of a neuregulin-receptor complex, although neuregulins do not interact with it alone. GP30 is a potential ligand for this receptor. Not activated by EGF, TGF-a and amphiregulin.

Product Info

Amount: 50 μl / 100 μl

Content: Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM

NaCl, 0.02% sodium azide and 50% glycerol.

Storage condition:

Storage condition:

Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid

repeated freeze and thaw cycles.

Application Note

Predicted MW: 185kd, Western blotting: 1:500~1:1000, Immunohistochemistry: 1:50~1:100, Immunofluorescence: 1:100~1:200




Figure 1: Western blot analysis of extracts from A431 cells untreated or treated with EGF using HER2(Phospho-Tyr877) Antibody 35-1073.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 2: Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue, using HER2(Phospho-Tyr877) Antibody 35-1073 (left) or the same antibody preincubated with blocking peptide(right).




Figure 3: Immunofluorescence staining of methanol-fixed MCF7 cells using HER2(Phospho-Tyr877) Antibody 35-1073.