

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 35-1078: Polyclonal Antibody to EGFR (Phospho-Ser1070)

Clonality: Polyclonal Application: WB,IHC Reactivity: Human **EGFR** Gene: Gene ID: 1956 P00533 **Uniprot ID:** Format: Purified

**Alternative Name:** ERBB1, Receptor protein-tyrosine kinase ErbB-1, kinase EGFR

Isotype: Rabbit IgG

Peptide sequence around phosphorylation site of serine 1070 (R-Y-S(p)-S-D) derived from Human Immunogen Information:

EGFR.

## **Description**

Receptor for EGF, but also for other members of the EGF family, as TGF-a, amphiregulin, betacellulin, heparin-binding EGF-like growth factor, GP30 and vaccinia virus growth factor. Is involved in the control of cell growth and differentiation. Phosphorylates MUC1 in breast cancer cells and increases the interaction of MUC1 with SRC and CTNNB1/beta-catenin.

## **Product Info**

 $50 \mu l / 100 \mu l$ Amount:

Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM Content:

NaCl, 0.02% sodium azide and 50% glycerol.

Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid Storage condition:

repeated freeze and thaw cycles.

## **Application Note**

Predicted MW: 175kd, Western blotting: 1:500~1:1000, Immunohistochemistry: 1:50~1:100




Figure 1: Western blot analysis of extracts from HUVEC cells untreated or treated with EGF using EGFR(Phospho-Ser1070) Antibody 35-1078.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 2: Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using EGFR(Phospho-Ser1070) Antibody 35-1078 (left) or the same antibody preincubated with blocking peptide(right).