

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

35-1124: Polyclonal Antibody to HDAC8 (Phospho-Ser39)

Clonality: Polyclonal
Application: IHC,IF
Reactivity: Human
Gene: HDAC8
Gene ID: 55869
Uniprot ID: Q9BY41
Format: Purified

Alternative Name: HD8, HDA8, HDACL1

Isotype: Rabbit IgG

Immunogen Information : Peptide sequence around phosphorylation site of serine 39 (R-A-S(p)-M-V) derived from Human

HDAC8.

Description

Histones play a critical role in transcriptional regulation, cell cycle progression, and developmental events. Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein encoded by HDAC8 gene belongs to class I of the histone deacetylase family. It catalyzes the deacetylation of lysine residues in the histone N-terminal tails and represses transcription in large multiprotein complexes with transcriptional co-repressors. Multiple transcript variants encoding different isoforms have been found for this gene.

Product Info

Amount: 50 μl / 100 μl

Content: Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM

NaCl, 0.02% sodium azide and 50% glycerol.

Storage condition:

Storage condition:

Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid

repeated freeze and thaw cycles.

Application Note

Predicted MW: 42kd, Immunohistochemistry: 1:50~1:100, Immunofluorescence: 1:100~1:200




Figure 1: Immunohistochemical analysis of paraffin-embedded human lung carcinoma tissue using HDAC8(Phospho-Ser39) Antibody 35-1124 (left) or the same antibody preincubated with blocking peptide(right).



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 2: Immunofluorescence staining of methanol-fixed Hela cells using HDAC8(Phospho-Ser39) Antibody 35-1124.