

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

35-1315: Polyclonal Antibody to PLC- Gamma2 (phospho-Tyr1217)

Clonality: Polyclonal

Application: IHC

Reactivity: Human, Mouse, Rat

 Gene :
 PLCG2

 Gene ID :
 5336

 Uniprot ID :
 P16885

 Format :
 Purified

Alternative Name: PLC-IV, PLC-gamma2, Phospholipase C-gamma-2

Isotype: Rabbit IgG

Immunogen Information: Peptide sequence around phosphorylation site of tyrosine 1217 (F-L-Y(p)-D-T) derived from

Human PLCg2.

Description

The production of the second messenger molecules diacylglycerol (DAG) and inositol 1,4,5-trisphosphate (IP3) is mediated by activated phosphatidylinositol-specific phospholipase C enzymes. It is a crucial enzyme in transmembrane signaling.

Product Info

Amount: 50 μl / 100 μl

Content: Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM

NaCl, 0.02% sodium azide and 50% glycerol.

Storage condition:

Storage condition:

Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid

repeated freeze and thaw cycles.

Application Note

Predicted MW: 150kd, Immunohistochemistry: 1:50~1:100




Figure 1: Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using PLC-g2(Phospho-Tyr1217) Antibody 35-1315 (left) or the same antibody preincubated with blocking peptide(right).