

35-1530: Polyclonal Antibody to AMPK Alpha1/AMPK Alpha2 (Ab-174/172)

Clonality :	Polyclonal
Application :	WB,IHC,IF
Reactivity :	Human,Mouse,Rat
Gene :	PRKAA1
Gene ID :	5562
Uniprot ID :	Q13131
Format :	Purified
Alternative Name :	AMPK,AMPK α 1
Isotype :	Rabbit IgG
Immunogen Information :	Peptide sequence around aa. 172~176/170~174 (L-R-T-S-C) derived from Human AMPK α 1/AMPK α 2.

Description

Responsible for the regulation of fatty acid synthesis by phosphorylation of acetyl-CoA carboxylase. It also regulates cholesterol synthesis via phosphorylation and inactivation of hormone-sensitive lipase and hydroxymethylglutaryl-CoA reductase. Appears to act as a metabolic stress-sensing protein kinase switching off biosynthetic pathways when cellular ATP levels are depleted and when 5'-AMP rises in response to fuel limitation and/or hypoxia. This is a catalytic subunit.

Product Info

Amount :	50 μ l / 100 μ l
Content :	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg $^{2+}$ and Ca $^{2+}$), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage condition :	Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid repeated freeze and thaw cycles.

Application Note

Predicted MW: 63kd, Western blotting: 1:500~1:1000, Immunohistochemistry: 1:50~1:100, Immunofluorescence: 1:100~1:200




Figure 1: Western blot analysis of extracts from A549, HepG2 and 3T3 cells using AMPK α 1/AMPK α 2(Ab-174/172) Antibody 35-1530 .




Figure 2: Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using AMPK α 1/AMPK α 2(Ab-174/172) Antibody 35-1530 (left) or the same antibody preincubated with blocking peptide(right).




Figure 3: Immunofluorescence staining of methanol-fixed HeLa cells using AMPK α 1/AMPK α 2(Ab-174/172) Antibody 35-1530 .