

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

35-1832: Polyclonal Antibody to GFI1b

Clonality: Polyclonal

Application: WB

Reactivity: Human, Mouse, Rat

 Gene :
 GFI1B

 Gene ID :
 8328

 Uniprot ID :
 Q5VTD9

 Format :
 Purified

 Alternative Name :
 Gfi-1b

 Isotype :
 Rabbit IgG

Immunogen Information: Peptide sequence around aa.322~326(H-R-E-S-Q) derived from Human GFI1b.

Description

Essential proto-oncogenic transcriptional regulator necessary for development and differentiation of erythroid and megakaryocytic lineages. Transcriptional repressor or activator depending on both promoter and cell type context; represses promoter activity of SOCS1 and SOCS3 and thus, may regulate cytokine signaling pathways. Cooperates with GATA1 to repress target gene transcription, such as the apoptosis regulator BCL2L1; GFI1B silencing in leukemic cell lines markedly increase apoptosis rate. Inhibits down-regulation of MYC and MYB as well as the cyclin-dependent kinase inhibitor CDKN1A/P21WAF1 in IL6-treated myelomonocytic cells. Represses expression of GATA3 in T-cell lymphomas and inhibits GATA1-mediated transcription; as GATA1 also mediates erythroid GFI1B transcription, both GATA1 and GFI1B participate in a feedback regulatory pathway controlling the expression of GFI1B gene in erythroid cells. Suppresses GATA1-mediated stimulation of GFI1B promoter through protein interaction. Binds to gamma-satellite DNA and to its own promoter, auto-repressing its own expression. Alters histone methylation by recruiting histone methyltransferase to target genes promoters. Plays a role in heterochromatin formation. Osawa M., Yamaguchi T., Nakamura Y. Iwama A.Blood 100:2769-2777(2002) Huang D.-Y., Kuo Y.-Y., Chang Z.-F. Nucleic Acids Res. 33:5331-5342(2005) Vassen L., Fiolka K., Moeroey T. EMBO J. 25:2409-2419(2006)

Product Info

Amount : 50 μl / 100 μl

Content: Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM

NaCl, 0.02% sodium azide and 50% glycerol.

Storage condition:

Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid

repeated freeze and thaw cycles.

Application Note

Predicted MW: 38kd, Western blotting: 1:500~1:1000



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 1: Western blot analysis of extracts from Raji cells using GFI1b Antibody 35-1832 .