

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

10-7543: Monoclonal Antibody to Arginase-1 (Clone: ABM4B35)

Clonality: Monoclonal ABM4B35 Clone Name: Application: IHC.WB Reactivity: Human Gene: ARG1 Gene ID: 383 **Uniprot ID:** P05089 Format: Purified **Alternative Name:** ARG1

Isotype: Mouse IgG2b Kappa

Immunogen Information: A partial length recombinant Arginase-1 protein (amino acids 11-230) was used as the immunogen

for this antibody

Description

Arginase-1 (Arg-1) is a binuclear manganese metalloenzyme that catalyzes the hydrolysis of arginine to ornithine and urea. It is expressed in normal human liver with a high degree of specificity, concentrated in periportal hepatocytes. Arginase-1 are urea cycle enzymes used to distinguish hepatocellular carcinoma from other carcinomas. Two transcript variants encoding different isoforms have been found for this gene. Inherited deficiency of this enzyme results in argininemia, an autosomal recessive disorder characterized by hyperammonemia.

Product Info

Amount : 25 μg / 100 μg

Purification: Protein G Chromatography

Content: 25 µg in 50 µl/100 µg in 200 µl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium

azide is highly toxic.

Storage condition:

Storage condition:

Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid

repeated freeze and thaw cycles.

Application Note

Western blot analysis: 2-4 µg/ml, Immunohistochemical analysis: 5-15 µg/ml




Fig-1: Immunohistochemical analysis of Arginase-1 $\,$ in human liver tissue using Arginase-1 antibody (Clone: ABM4B35) at 15 μ g/ml.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Fig-2 : Immunohistochemical analysis of Arginase-1 in human liver tissue using Arginase-1 antibody (Clone: ABM4B35) at 5 $\mu g/ml$.




Fig-3 : Western blot analysis of Arginase-1. Anti- Arginase-1 antibody (Clone: ABM4B35) was tested at 2 μ g/ml on human liver lysate.