

## 32-12058: Human Fibroblast Growth Factor-10

**Gene :** FGF10  
**Gene ID :** 2255  
**Uniprot ID :** O15520  
**Alternative Name :** Keratinocyte growth factor 2

### Description

**Source:** Genetically modified E.coli.

**Predicted MW:** Monomer, 19.3 kDa (170 aa)

Fibroblast growth factor 10 (FGF-10) is a growth factor that is important during embryonic development, especially during lung, limb, brain, heart, and kidney morphogenesis. FGF-10 is expressed in mesenchymal cells and facilitates epithelial-mesenchymal signaling through binding the epithelially expressed FGF receptor 2b (FGFR2b). FGF-10 also functions as a mitogen for keratinizing epidermal cells, and induces the migration and invasion of cancer cells.

### Product Info

**Amount :** 25 µg / 100 µg  
**Purification :** Reducing and Non-Reducing SDS PAGE at  $\geq 95\%$   
Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 10 mM sodium phosphate, pH 7.5  
Sterile water at 0.1 mg/mL  
**Content :**  
**Storage condition :** Store at  $-20^{\circ}\text{C}$   
**Amino Acid :** MLGQDMVSPE ATNSSSSSFS SPSSAGRHVR SYNHLQGDVR WRKLSFTKY FLKIEKNGKV SG  
TKKENCPY SILEITSVEIGVVAVKAINS NYLAMNKKG KLYGSKEFNN DCKLKERIEE NGYNTYAS  
FN WQHNGRQMYV ALNGKGAPRR GQKTRRKNTS AHFLPMVVHS

### Application Note

**Endotoxin:** Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.

Biological Activity was determined by 4MBR-5 cell proliferation at  $\leq 200$  ng/mL;  $\geq 5.0 \times 10^3$  units/mg (typical ED50 is  $< 80$  ng/mL). Centrifuge vial before opening, Suspend the product by gently pipetting the above recommended solution down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution. For prolonged storage, dilute to working aliquots in a 0.1% BSA solution, store at  $-80^{\circ}\text{C}$  and avoid repeat freeze thaws. Upon reconstitution, a small amount of visible precipitate can be expected. A 10% overfill has been added to the total material vialled to compensate for this loss.



