

## 36-3372: Anti-von Willebrand Factor / Factor VIII Related-Ag (Endothelial Marker) Monoclonal Antibody(Clone: F8/86)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	F8/86
<b>Application :</b>	IHC
<b>Reactivity :</b>	Human
<b>Gene :</b>	VWF
<b>Gene ID :</b>	7450
<b>Uniprot ID :</b>	P04275
<b>Alternative Name :</b>	Coagulation Factor VIII, Factor VIII Related Antigen, F8VWF, von Willebrand Antigen 2, von Willebrand Disease (vWD)
<b>Isotype :</b>	Mouse IgG1, kappa
<b>Immunogen Information :</b>	von Willebrand Factor isolated from human plasma

### Description

von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposi's sarcoma and cardiac myxoma. It is widely used for differentiating vascular lesions from those of other tissue differentiation within a panel of other vascular markers although not all tumors of endothelial differentiation contain this antigen.

### Product Info

<b>Amount :</b>	20 µg / 100 µg
<b>Content :</b>	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
<b>Storage condition :</b>	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

### Application Note

Immunohistochemistry (Formalin-fixed) (0.1-0.2ug/ml for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95&degC followed by cooling at RT for 20 minutes);




Fig. 1: Formalin-fixed, paraffin-embedded human Tonsil stained with vWF Mouse Monoclonal Antibody (F8/86).




Fig. 2: SDS-PAGE Analysis Purified vWF Mouse Monoclonal Antibody (F8/86).  
Confirmation of Purity and Integrity of Antibody.