

### 36-3414: Anti-Eosinophil Peroxidase (EPX) Monoclonal Antibody(Clone: AHE-1)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	AHE-1
<b>Application :</b>	ELISA
<b>Reactivity :</b>	Human
<b>Gene :</b>	EPX
<b>Gene ID :</b>	8288
<b>Uniprot ID :</b>	P11678
<b>Alternative Name :</b>	Eosinophil peroxidase heavy chain; EPER; EPO; EPP; EPX; EPX PEN; PERE
<b>Isotype :</b>	Mouse IgG1, kappa
<b>Immunogen Information :</b>	Human eosinophils from a patient with hypereosinophilic syndrome

#### Description

Peripheral blood granulocytes are classified into neutrophils, basophils and eosinophils according to the staining characteristics of their cytoplasmic granules. Granule proteins are released by physiologic and pharmacologic stimuli and play important roles in both normal and pathological host immune responses. Eosinophil major basic protein and eosinophil peroxidase (EPX) are granule proteins specific to the eosinophil. AHE-1 recognizes human EPX, a granule protein specific to eosinophils. It does not cross-react with eosinophil major basic protein, elastase, cathepsin G, esterase N, thrombin, plasmin, kallikrein, lactoferrin, or transferrin. This MAb stains eosinophils only and does not stain other peripheral blood cells, including platelets, neutrophils, monocytes, lymphocytes or red blood cells. Human EPX gene product can form a tetramer of two light chains and two heavy chains. Other peroxidase family members include myeloperoxidase (MPO), lactoperoxidase (LPO), and thyroid peroxidase (TPO).

#### Product Info

<b>Amount :</b>	20 µg / 100 µg
<b>Content :</b>	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
<b>Storage condition :</b>	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

#### Application Note

ELISA (For coating, order antibody without BSA);,