

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

36-3688: Anti-GITR / Tnfrsf18 (Mouse) Monoclonal Antibody(Clone: DTA-1)

Clone Name: DTA-1
Application: FACS,IF
Reactivity: Mouse
Gene ID: 21936
Uniprot ID: O35714

Activation-inducible TNF-related ligand; AITRL; GITRL; Glucocorticoid induced TNFR related

Alternative Name: protein ligand; Glucocorticoid-induced TNF-related ligand; hGITRL; TL6; TNFSF18; Tumor

necrosis factor (ligand) superfamily member 18

Isotype: Rat IgG2b, lambda

Immunogen Information: Mouse CD25+ CD4+ T cells

Description

GITR (glucocorticoid-induced TNFR-related gene) is a member of the TNF-receptor superfamily, also known as TNFRSF18. It is expressed at low levels on resting T lymphocytes and at high levels on CD25+CD4+Tregs. The expression of GITR on T cells can be upregulated upon activation. Interaction of GITR with its ligand (GITRL) has been demonstrated to aµgment T cell activation, proliferation, cytokine production as well as MAPKs and NF-B activation, and abrogate the inhibitory function of CD25+CD4+Tregs. In vivoactivation of GITR causes development of autoimmune diseases and restores the suppressed immune response.

Product Info

Amount : $20 \mu g / 100 \mu g$

Content: 200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with

0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage condition:

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is

stable for 24 months. Non-hazardous.

Application Note

Flow Cytometry (0.5-1ug/million cells in 0.1ml); Immunofluorescence (1-2ug/ml);




Fig. 1: SDS-PAGE Analysis Purified GITR Mouse Monoclonal Antibody (DTA-1). Confirmation of Purity and Integrity of Antibody