

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

36-3821: Anti-CD40, Mouse Monoclonal Antibody(Clone: FGK45.5)

Clonality: Monoclonal Clone Name: FGK45.5

Application: Functional Assay, FACS, IF

 Reactivity :
 Mouse

 Gene ID :
 21939

 Uniprot ID :
 P27512

Alternative Name:

B cell surface antigen CD40, CD40 type II isoform, p50, Bp50, CD40L receptor, CDW40, TRAP,

IMD3, IGM, Tnfrsf5, HIGM1, GP39, T-BAM

Isotype: Rat IgG2a, kappa

Immunogen Information: Purified recombinant Mouse CD40 fusion protein

Description

Upon binding to its ligand CD154, CD40 acts as a costimulatory molecule for the activation of B cells, dendritic cells, monocytes, and other APCs. CD40 plays roles in B cell activation, differentiation, proliferation and Ig isotype switching as well as dendritic cell maturation. FGK4.5 can be used for blocking of CD40/CD154 interaction and forin vitroandin vivoactivation of CD40 expressing APCs. This MAb is shown to indirectly activate natural killer (NK) cells, producing significant antitumor and antimetastatic effects. Effective in boosting immune responses against infectious agents and can potentially be used to treat chronic autoimmune inflammatory processes.

Product Info

Amount : 20 μg / 100 μg

Content: 200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with

0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage condition:

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is

stable for 24 months. Non-hazardous.

Application Note

Functional Studies (Order Antibody without BSA & Azide); Flow Cytometry (0.5-1ug/million cells in 0.1ml); Immunofluorescence (1-2ug/ml);