

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-2331: FTCD Recombinant Protein

Alternative Name: Formiminotransferase-cyclodeaminase, FTCD, LCHC1, LC-1.

Description

Source: Sf9 insect cells. Formiminotransferase Cyclodeaminase Human Recombinant (also called liver cytosol type 1) produced in SF9, is a glycosylated, polypeptide chain having a molecular mass of 59,749 Dalton. The FTCD is expressed with a -6xHis tag and purified by proprietary chromatographic techniques. Formiminotransferase cyclodeaminase is an enzyme which catalyzes the conversion of formiminoglutamate and tetrahydrofolate into formiminotetrahydrofolateand glutamate. Formiminotransferase cyclodeaminase, a bifunctional enzyme of tetrahydrofolate synthesis, is the target antigen of anti-LC1 (liver cytosol antigen type 1) autoantibodies. Presence of LC1 autoantibodies is a marker for type 2 autoimmune hepatitis (for which anti-LKM 1/cytochrome P450 2D6 autoantibodies are a further marker). Serum LC1 autoantibody concentrations appear to fluctuate in parallel with aminotransferase levels, a particularly intriguing observation that suggests a possible role of LC1 autoreactivity in the pathogenic mechanism leading to hepatocyte injury.

Product Info

Amount: 10 µg

Purification: Greater than 95% as determined by SDS-PAGE.

Content: FTCD is supplied in 16mM HEPES buffer pH-7.6, 240mM sodium chloride, and 20% glycerol.

Storage condition:

Storage condition:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°Cfor longer periods of

time. Avoid multiple freeze-thaw cycles.

