

36-1309: Monoclonal Antibody to Insulin / IRDN (beta-Cell & Insulinoma Marker)(Clone : SPM139)

Clonality :	Monoclonal
Clone Name :	SPM139
Application :	FACS,IF,IHC
Reactivity :	Human
Gene :	INS
Gene ID :	3630
Uniprot ID :	P01308
Format :	Purified
Alternative Name :	INS
Isotype :	Mouse IgG1, kappa
Immunogen Information :	Full length (1-84 amino acid) purified pig insulin, conjugated to KLH

Description


Recognizes a polypeptide which is identified as insulin, a 51-amino acid polypeptide composed of A and B chains connected through the C-peptide. Proinsulin, which has very little biological activity, is cleaved by proteases within its cell of origin into the insulin molecule and the C-terminal basic residue. Insulin enhances membrane transport of glucose, amino acids, and certain ions. It also promotes glycogen storage, formation of triglycerides, and synthesis of proteins and nucleic acids. Deficiency of insulin results in diabetes mellitus. The main storage site for insulin is the pancreatic islets. Antibodies to insulin are important as beta-cell and insulinoma marker.

Product Info

Amount :	100 µg
Purification :	Affinity Chromatography
Content :	100 µg in 500 µl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly toxic.
Storage condition :	Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid repeated freeze and thaw cycles.

Application Note

Flow Cytometry (1-2ug/million cells); Immunofluorescence (1-2ug/ml); Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(No special pretreatment is required for staining of formalin/paraffin tissues.)



Formalin-fixed, paraffin-embedded human Pancreas stained with Insulin Monoclonal Antibody (SPM139).