

### 36-1342: Monoclonal Antibody to Cytokeratin 6 (KRT6) (Hyperproliferation-Related Keratin)(Clone : LHK6; same as LHK6B)(Discontinued)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	LHK6; same as LHK6B
<b>Application :</b>	FACS,IF,IHC
<b>Reactivity :</b>	Human,Mouse
<b>Gene :</b>	KRT6A
<b>Gene ID :</b>	3853
<b>Uniprot ID :</b>	P02538
<b>Format :</b>	Purified
<b>Alternative Name :</b>	KRT6A,K6A,KRT6D
<b>Isotype :</b>	Mouse IgG2a
<b>Immunogen Information :</b>	A synthetic peptide of 11 amino acids (GSSTIKYTTTS) from C-terminus of human keRatin 6

#### Description

This MAb recognizes a protein of 56kDa, identified as cytokeratin 6 (CK6). In humans, multiple isoforms of Cytokeratin 6 (6A-6F), encoded by several highly homologous genes, have distinct tissue expression patterns, and Cytokeratin 6A is the dominant form in epithelial tissue. The gene encoding human Cytokeratin 6A maps to chromosome 12q13, and mutations in this gene are linked to several inheritable hair and skin pathologies. Keratins 6 and 16 are expressed in keratinocytes, which are undergoing rapid turnover in the suprabasal region (also known as hyper-proliferation-related keratins). Keratin 6 is found in hair follicles, suprabasal cells of a variety of internal stratified epithelia, in epidermis, in both normal and hyper-proliferative situations. Epidermal injury results in activation of keratinocytes, which express CK6 and CK16. CK6 is strongly expressed in about 75% of head and neck squamous cell carcinomas. Expression of CK6 is particularly associated with differentiation.


#### Product Info

<b>Amount :</b>	100 µg
<b>Purification :</b>	Affinity Chromatography
<b>Content :</b>	100 µg in 500 µl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly toxic.
<b>Storage condition :</b>	Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid repeated freeze and thaw cycles.

#### Application Note

Flow Cytometry (0.5-1.0 µg/million cells in 0.1ml); Immunofluorescence (0.5-1.0 µg/ml); Immunohistology (Formalin-fixed) (0.1-0.2 µg/ml for 30 min at RT); (Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes); Optimal dilution for a specific application should be determined.





Formalin-fixed, paraffin-embedded human Bladder Carcinoma stained with Cytokeratin 6 Monoclonal Antibody (LHK6)