

32-2015: Deslorelin

Description

Deslorelin is a potent LHRH/GnRH agonist has a molecular formula of $C_{64}H_{83}N_{17}O_{12}$, Pyr-His-Trp-Ser-Tyr-D-Trp-Leu-Arg-Pro-NH₂ having an Mw of 1281.64 Dalton. Deslorelin is being studied in the treatment of cancer as a way to block sex hormones made by the ovaries or testicles. It belongs to the family of drugs called gonadotropin-releasing hormone analogs. It is used for the induction of ovulation in mares. Deslorelin binds to and activates pituitary gonadotropin releasing hormone (GnRH) receptors. Continuous, prolonged administration of goserelin in males results in pituitary GnRH receptor desensitization and inhibition of pituitary secretion of follicle stimulating hormone (FSH) and luteinizing hormone (LH), leading to a significant decline in testosterone production; in females, prolonged administration results in a decrease in estradiol production.

Product Info

| | |
|----------------------------|--|
| Amount : | 25 mg |
| Purification : | Greater than 99.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE. |
| Content : | The Deslorelin peptide was lyophilized with no additives. |
| Storage condition : | Lyophilized Deslorelin although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution Deslorelin should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Please prevent freeze-thaw cycles. |

Application Note

It is recommended to reconstitute the lyophilized Deslorelin in sterile 18M Ω -cm H₂O not less than 100 μ g/ml, which can then be further diluted to other aqueous solutions.

