9853 Pacific Heights Blvd.
Suite D. San Diego,
CA 92121, USA
Tel: 858-263-4982
Email: info@abeomics.com

32-2094: Vasopressin Protein

Alternative Name :	Vasopressin-neurophysin 2-copeptin,AVP-NPII,VP,ADH,ARVP,AVRP,Argipressin,Arginine vasopressin,AVP,Vasopressin,Antidiuretic Hormone.

Description

8-L-Arginine Vasopressin's molecular weight is 1084.24 Dalton and the molecular formula is: C46H65N15O12S2. Arginine vasopressin (AVP), also known as argipressin or antidiuretic hormone (ADH), is a human hormonethat is released when the body is low on water; it causes the kidneysto conserve water, but not salt, by concentrating the urineand reducing urine volume. It also raises blood pressure by inducing moderate vasoconstriction. It has various effects in the brain.A very similar substance, lysine vasopressin (LVP) or lypressin, has the same function in pigsand is often used in human therapy.Vasopressin is a peptide hormone. It is derived from a preprohormoneprecursor that is synthesized in the hypothalamus, from which it is liberated during transport to the posterior pituitary. Most of it is stored in the posterior partof the pituitary glandto be released into the blood stream; some of it is also released directly into the brain.AVP allows water reabsorption by the of additional water channels in cortical and inner medullary collecting ducts.

Product Info

Amount :	25 mg
Purification :	Greater than 98.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.
Content :	The protein (1 mg/ml) was lyophilized with no additives.
Storage condition :	Lyophilized Vasopressin although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution Vasopressin should be stored at 4°C between 2-7 days and for future use below -18°C.For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Please prevent freeze-thaw cycles.
Amino Acid :	C[Cys-Tyr-Phe-Gln-Asn-Cys]-Pro-Arg-Gly-NH2.

Application Note

It is recommended to reconstitute the lyophilized Vasopressin in sterile 18MÃ□Â©-cm H2O not less than 100 Ã□Âµg/ml, which can then be further diluted to other aqueous solutions.

For Research Use Only. Not for use in diagnostic/therapeutics procedures.