

32-2242: Cyclophilin B Recombinant Protein

Alternative Name : Peptidylprolyl isomerase B,PPIase,Rotamase,S-cyclophilin,PPIB,cyclophilin-like protein,peptidyl-prolyl cis-trans isomerase B,Cyclophilin B,SCYLP,CYPB,CYP-S1,MGC2224,MGC14109.

Description

Source : Escherichia Coli. Cyclophilin-B Human Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 192 amino acids (26-216) and having a molecular mass of 21.2 kDa. PPIB is purified by proprietary chromatographic techniques. Cyclophilin B (also known as PPIB, peptidylpropyl isomerase B) is a cyclosporine-binding protein and is mainly located within the endoplasmic reticulum. It is associated with the secretory pathway and released in biological fluids. This protein can bind to cells derived from T- and B-lymphocytes, and may regulate cyclosporine A-mediated immunosuppression.

Product Info

Amount : 50 µg
Purification : Greater than 95.0% as determined by SDS-PAGE.
Content : 1 mg/ml solution containing 20mM Tris-HCl 8.0, 20mM NaCl, 0.5mM DTT and 10% glycerol.
Storage condition : Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.
Amino Acid : MLLPGPSAAD EKKKGPKVTV KVFYDLRIGD EDVGRVIFGL FGKTVPKTVD NFVALATGEKGFYKNSKFH
 RVIKDFMIQG GDFTRGDGTG GKSIYGERFP DENFKLKHYG PGWVSMANAGKDTNGSQFFI TTVKTAWLDG
 KHVVFQKVL EMEVVRKVES TKTDSDRKPL KDVIADCGK IEVEKPFAIA KE.

Application Note

Specific activity is > 220 nmoles/min/mg, and is defined as the amount of enzyme that cleaves 1umole of suc-AAFP-pNA per minute at 25C in Tris-Hcl pH8.0 using chymotrypsin.

