9853 Pacific Heights Blvd.
Suite D. San Diego,
CA 92121, USA
Tel: 858-263-4982
Email: info@abeomics.com

32-2617: NTH Recombinant Protein

Alternative Name : DNA-(apurinic or apyrimidinic site) lyase,b1633,JW1625.

Description

Source : Escherichia Coli. NTH E.Coli Recombinant fused with a 20 amino acid His tag at N-terminus produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 231 amino acids (1-211a.a.) and having a molecular mass of 25.7kDa. The NTH is purified by proprietary chromatographic techniques. Endonuclease III (nth) is a DNA repair enzyme which has both DNA N-glycosylase activity and AP-lyase activity. The DNA N-glycosylase activity releases numerous damaged pyrimidines from DNA by cleaving the N-glycosidic bond and leaving an AP (apurinic/apyrimidinic) site. This AP-lyase activity cleaves the phosphodiester bond 3' to the AP site by a beta-elimination, thus leaving a 3'-terminal unsaturated sugar and a product with a terminal 5'-phosphate.

Product Info

Amount :	10 µg
Purification :	Greater than 85.0% as determined by SDS-PAGE.
Content :	The NTH solution (0.5mg/1ml) contains 20mM Tris-HCl buffer (pH8.0), 0.1M NaCl, 1mM DTT, 0.1mM PMSF and 40% glycerol.
Storage condition :	Store at 4°C if entire vial will be used within 2-4 weeks.Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Avoid multiple freeze-thaw cycles.
Amino Acid :	MGSSHHHHHH SSGLVPRGSH MNKAKRLEIL TRLRENNPHP TTELNFSSPF ELLIAVLLSA QATDVSVNKA TAKLYPVANT PAAMLELGVE GVKTYIKTIG LYNSKAENII KTCRILLEQH NGEVPEDRAA LEALPGVGRK TANVVLNTAF GWPTIAVDTH IFRVCNRTQF APGKNVEQVE EKLLKVVPAE FKVDCHHWLI LHGRYTCIAR KPRCGSCIIE DLCEYKEKVD I.

For Research Use Only. Not for use in diagnostic/therapeutics procedures.