

32-2680: PLA2G2D Recombinant Protein

Alternative Name : Group IID secretory phospholipase A2, EC 3.1.1.4, Phosphatidylcholine 2-acylhydrolase GIID, GIID sPLA2, PLA2IID, sPLA(2)-IID, Secretory-type PLA, stroma-associated homolog, SPLASH, sPLA2S, PLA2G2D.

Description

Source : Escherichia Coli. Secreted Phospholipase A2-IID Human Recombinant was produced with N-terminal His-Tag. PLA2G2D His-Tagged Fusion protein is 16.4 kDa containing 125 amino acid residues of the human secreted phospholipase A2-IID and 16 additional amino acid residues - His-Tag . Phospholipase A2 (PLA2) catalyzes the hydrolysis of the sn-2 position of membrane glycerophospholipids to liberate arachidonic acid (AA), a precursor of eicosanoids including prostaglandins and leukotrienes. The same reaction also produces lysophospholipids, which represent another class of lipid mediators. The secretory PLA2 (sPLA2) family, in which 10 isozymes have been identified, consists of low molecular weight, Ca²⁺-requiring secretory enzymes that have been implicated in a number of biological processes, such as modification of eicosanoid generation, inflammation, and host defense. This enzyme has been proposed to hydrolyze phosphatidylcholine (PC) in lipoproteins to liberate lyso-PC and free fatty acids in the arterial wall, thereby facilitating the accumulation of bioactive lipids and modified lipoproteins in atherosclerotic foci. In mice, sPLA2 expression significantly influences HDL particle size and composition and demonstrate that an induction of sPLA2 is required for the decrease in plasma HDL cholesterol in response to inflammatory stimuli. Instillation of bacteria into the bronchi was associated with surfactant degradation and a decrease in large: small ratio of surfactant aggregates in rats.

Product Info

Amount : 10 µg
Content : Sterile filtered and lyophilized from 0.5 mg/ml in 0.05M Acetate buffer pH-4.
Storage condition : Store lyophilized protein at -20°C. Aliquot the product after reconstitution to avoid repeated freezing/thawing cycles. Reconstituted protein can be stored at 4°C for a limited period of time; it does not show any change after two weeks at 4°C.
Amino Acid : MRGSHHHHHHGMASHMGILNLNKMVKQVTGKMPILSYWPYGCHCGLGGR
GQPKDATDWCCQTHDCCYDHLKTQGCGIYKYRYNFSQGNIHCSDKGSWC
EQQLCADCKEVAFCLKRNLDTYQRLRFYWRPHCRGQTPGC.

Application Note

Add 0.2 ml of 0.1M Acetate buffer pH-4 and let the lyophilized pellet dissolve completely. For conversion into higher pH value, we recommend intensive dilution by relevant buffer to a concentration of 10 µg/ml. In higher concentrations the solubility of this antigen is limited.

