

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-3058: MEK1 Recombinant Protein

Alternative Name MEK1,MKK1,MAPKK1,MAPZK1,Dual specificity mitogen-activated protein kinase kinase 1,MAP kinase kinase 1,EC 2.7.12.2,ERK activator kinase 1,MAPK/ERK kinase 1,PRKMK1.

Description

Source: Sf9 Insect Cells. MAP2K1 active Human Recombinant produced in Sf9 cells is a glycosylated, polypeptide chain containing amino acids 2-393 having a molecular mass of 47 kDa. MAP2K1 is fused to a polyhistidine tag and is purified by proprietary chromatographic techniques. MAP2K1 is a member of the dual specificity protein kinase family, which plays a role as a mitogen-activated protein (MAP) kinase kinase. MAP kinases, are recognized as extracellular signal-regulated kinases (ERKs), that act as an integration position for multiple biochemical signals. MEK1 is located upstream of MAP kinases and stimulates the enzymatic activity of MAP kinases upon wide variety of extra- and intracellular signals. As a key player of MAP kinase signal transduction pathway, MEK1 is involved in many cellular processes such as proliferation, differentiation, transcription regulation and development. MAP2K1 catalyzes the concomitant phosphorylation of a threonine and a tyrosine residue in a thr-glu-tyr sequence located in map kinases. MEK1 activates erk1 and erk2 map kinases.

Product Info

Content:

Amount: $5 \mu g$

Purification: Greater than 90% as determined by SDS-PAGE.

MEK1 is supplied at a concentration of 0.09mg/ml in 40mM Tris, pH-8, 0.15M NaCl, 0.27M

sucrose, 1mM DTT, 0.2mM PMSF, 1mM benzamidine, 0.1mM sodium vanadate and 0.03%

Brij-35.

Storage condition:

Store at 4°C if entire vial will be used within 1-2 weeks. Store frozen at -20°C for longer periods of time. Avoid multiple frozen than evelop.

of time. Avoid multiple freeze-thaw cycles.

Application Note

 \sim 125-175 units/mg. One unit of MEK1 activity transfers 1 nmol of phosphate to ERK1/2 peptide per minute at $30\tilde{A} \square \hat{A}^{\circ} C$ in a reaction containing $100\tilde{A} \square \hat{A} \mu M$ ATP. Recombinant active MEK1 also phosphorylates ERK1, ERK2, and GSK-3?. Kinase activity may vary depending on the substrate and reaction conditions. The optimal concentration should be determined for each specific application.

