

32-3115: STK16 Recombinant Protein

Alternative Name : Serine/threonine-protein kinase 16, Myristoylated and palmitoylated serine/threonine-protein kinase, MPSK, Protein kinase PKL12, TGF-beta-stimulated factor 1, TSF-1, Tyrosine-protein kinase STK16, hPSK, STK16, MPSK1, PKL12, TSF1, KRCT.

Description

Source : Escherichia Coli. STK16 Human Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 329 amino acids (1-305 a.a) and having a molecular mass of 37.2kDa. STK16 is fused to a 24 amino acid His-tag at N-terminus & purified by proprietary chromatographic techniques. Serine/threonine-protein kinase 16 (STK16) is a membrane-associated protein kinase which phosphorylates on serine and threonine residues. STK16 is involved in secretory vesicle trafficking or intracellular signaling. Furthermore, the STK16 protein may have a role in regulating stromal-epithelial interactions which occur during ductal morphogenesis in the mammary gland. STK16 can autophosphorylate on Tyr residue; it is however unclear whether STK16 has tyrosine-protein kinase toward other proteins. STK16 may also be involved in TGF-beta signaling.

Product Info

Amount : 20 µg
Purification : Greater than 90.0% as determined by SDS-PAGE.
Content : STK16 protein solution (1mg/ml) containing 20mM Tris-HCl buffer (pH8.0), 20% glycerol, 0.1M NaCl and 1mM DTT.
Storage condition : Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.
Amino Acid : MGSSHHHHHH SSGLVPRGSH MGSHMGHALC VCSRGTVIID NKRYLFIQKL GEGGFSYVDL
 VEGLDHGHFY ALKRILCHEQ QDREEAQREA DMHRLFNHPN ILRLVAYCLR ERGAKHEAWL LLPFFKRGT
 WNEIERLKDK GNFLTEDQIL WLLGICRGL EAIHAKGYAH RDLKPTNILL GDEGQPVLM D LGSMNQACIH
 VEGSRQALT L QDWAAQRCTI SYRAPELFSV QSHCVIDERT DVWSLGC VLY AMMFGE GEPYD
 MVFQKGDSVA LAVQNQLSIP QSPRHSSALR QLLNSMMTV D PHQRPHIPLL LSQLEALQPP APGQHTTQI.

