

32-3301: BCL10 Recombinant Protein

Alternative Name :

B-cell CLL/lymphoma 10,B-cell lymphoma/leukemia 10,Mammalian CARD-containing adapter molecule E10,CARD-containing molecule enhancing NF-kappa-B,CED-3/ICH-1 prodomain homologous E10-like regulator,CARD-containing apoptotic signaling protein

Description

Source : E.coli. BCL10 Human Recombinant produced in E. coli is a single polypeptide chain containing 257 amino acids (1-233) and having a molecular mass of 28.8kDa (molecular weight on SDS-PAGE will appear higher).BCL10 is fused to a 24 amino acid His-tag at N-terminus & purified by proprietary chromatographic techniques. BCL10 holds a caspase recruitment domain (CARD), and is known as an inducer of apoptosis and as NF-kappaB activator. BCL10 cooperates with other CARD domain containing proteins such as CARD9, 10, 11 and 14, which operates as upstream regulators in NF-kappaB signaling. BCL10 is known to create a complex with MALT1, a protein encoded by another gene and is translocated in MALT lymphoma. MALT1 and BCL10 protein synergize the activation of NF-kappaB, and the deregulation of either of them facilitates the same pathogenetic process which leads to the malignancy.

Product Info

Amount :	5 µg
Purification :	Greater than 85% as determined by SDS-PAGE.
Content :	The BCL10 solution (1mg/ml) contains 20mM Tris-HCl buffer (pH 8.0), 50mM NaCl, 1mM DTT and 10% glycerol.
Storage condition :	Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Avoid multiple freeze-thaw cycles.
Amino Acid :	MGSSHHHHHH SSGLVPRGSH MGSHEPTAP SLTEEDLTEV KKDALENLRV YLCEKIAER HFDHLRAKKI LSREDTEEIS CRTSSRKRAK KLLDYLQENP KGLDTLVESI RREKTQNFLI QKITDEVLKL RNIKLEHLKG LKCSSEPFPP DGATNNLSRS NSDESNFSEK LRASTVMYHP EGESSTPPFF STNSSLNLPV LEVGRTEI FSSTTLPRPG DPGAPPLPPD LQLEEEGTCA NSSEMFLPLR SRTVSRQ.