

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-3320: Bcl-XL His Recombinant Protein

**Alternative Name:** BcIXL,BcI-X(L),BcI-XL.

## **Description**

Source: Escherichia Coli. Bcl-XL Human Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing amino acids 1-210. The Bcl-XL is expressed as His-Tag fusion protein and purified by proprietary chromatographic techniques. Bcl-XL is a transmembrane protein located in the mitochondrial membranes of cells that are long-lived and postmitotic, such as adult brain cells. It plays arole in the signal transduction pathway of the FAS-Ligand. Bcl-XL is an antiapoptotic protein which is a member of the Bcl-2 family which are able to form heterodimers, and this is an significant event in the regulation of apoptosis. BCL-XL is involved in the survival of cancer cells. Bcl-xL is the leading monitor of apoptosis/active cell suicide. Bcl-xL has cell death repressor activity and therefore acts as a survival protein.

## **Product Info**

Amount: 10 µg

**Purification:** Greater than 95.0% as determined by SDS-PAGE.

Content: The protein contains 10mM Tris-HCL pH-8, 1mM EDTA and 250mM NaCl.

Lyophilized Bcl-XL although stable at room temperature for 3 weeks, should be stored

desiccated below -18°C. Upon reconstitution Bcl-XL should be stored at 4°C between 2-7 days **Storage condition:** 

and for future use below -18°C.For long term storage it is recommended to add a carrier protein

(0.1% HSA or BSA). Please prevent freeze-thaw cycles.

## **Application Note**

Suspend BclXL in 100̸µl of 0.5M Acetic acid, over night at 4Ã∏°C. Dilute 10 fold into selected buffer system.Bcl-XL has tendency to form intramolecular disulfide bond, 5mM DTT is recommended in assay buffer. When running SDS-PAGE gel, 10mM DTT is recommended.

