

## 32-3455: sCD23 Recombinant Protein

**Alternative Name :** Low affinity immunoglobulin epsilon Fc receptor,Lymphocyte IgE receptor,Fc-epsilon-RII,BLAST-2,Immunoglobulin E-binding factor,CD23 antigen,FCER2,CD23,FCE2,CD23A,IGEBF,CLEC4J.

### Description

Source : Escherichia Coli. sCD23 Human Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 172 amino acids and having a molecular mass of 19.2kDa.The sCD23 is purified by proprietary chromatographic techniques. CD23 is a 45kDa glycoprotein, which is present on a subpopulation of freshly isolated peripheral blood and tonsil B cells and strongly expressed on EBV-transformed B lymphoblasts. The CD23 molecule is identical to the low affinity IgE receptor found on B cells. Expression of CD23 has been detected in neoplastic cells from cases of B cell chronic lymphocytic leukaemia and some cases of centroblastic/centrocytic lymphoma.

### Product Info

**Amount :** 20 µg  
**Purification :** Greater than 97.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.  
**Content :** Lyophilized from a 0.2µm filtered concentrated solution in PBS, pH 7.4.  
**Storage condition :** Lyophilized sCD23 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution sCD23 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.  
**Amino Acid :** MELQVSSGFV CNTCPEKWIN FQRKCYFYGK GTKQVWHARY ACDDMEGQLV SIHSPPEEQDF  
LTKHASHTGS WIGLRNLDLK GEFIWVDGSH VDYSNWAPGE PTSRSQGEDC VMMRGSGRWN  
DAFCDRKLGAVVCDRLATCT PPASEGSAES MGPDSRPPDP GRLPTPSAPL HS

### Application Note

It is recommended to reconstitute the lyophilized sCD23 in sterile 18M-cm H2O not less than 100µg/ml, which can then be further diluted to other aqueous solutions. The ED50 is determined by its ability to induce TNF-alpha production by human PBMCs.