

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-3455: sCD23 Recombinant Protein

Alternative Low affinity immunoglobulin epsilon Fc receptor, Lymphocyte IgE receptor, Fc-epsilon-

Name: RII,BLAST-2,Immunoglobulin E-binding factor,CD23 antigen,FCER2,CD23,FCE2,CD23A,IGEBF,CLEC4J.

Description

Source: Escherichia Coli. sCD23 Human Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 172 amino acids and having a molecular mass of 19.2kDa.The sCD23 is purified by proprietary chromatographic techniques. CD23 is a 45kDa glycoprotein, which is present on a subpopulation of freshly isolated peripheral blood and tonsil B cells and strongly expressed on EBV-transformed B lymphoblasts. The CD23 molecule is identical to the low affinity IgE receptor found on B cells. Expression of CD23 has been detected in neoplastic cells from cases of B cell chronic lymphocyctic leukaemia and some cases of centroblastic/centrocytic lymphoma.

Product Info

Amount: 20 µg

Purification: Greater than 97.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.

Content: Lyophilized from a 0.2µm filtered concentrated solution in PBS, pH 7.4.

Lyophilized sCD23 although stable at room temperature for 3 weeks, should be stored desiccated

below -18°C. Upon reconstitution sCD23 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or

BSA). Please prevent freeze-thaw cycles.

Amino Acid: MELQVSSGFV CNTCPEKWIN FQRKCYYFGK GTKQWVHARY ACDDMEGQLV SIHSPEEQDF

LTKHASHTGS WIGLRNLDLK GEFIWVDGSH VDYSNWAPGE PTSRSQGEDC VMMRGSGRWN

DAFCDRKLGA WVCDRLATCT PPASEGSAES MGPDSRPDPD GRLPTPSAPL HS

Application Note

Storage condition:

It is recommended to reconstitute the lyophilized sCD23 in sterile 18M-cm H2O not less than $100 \text{\AA} \mu\text{g/ml}$, which can then be further diluted to other aqueous solutions. The ED50 is determined by its ability to induce TNF-alpha production by human PBMCs.

