

## 32-3460: CD27 Recombinant Protein

**Alternative Name :** Tumor Necrosis Factor Receptor Superfamily Member 7,T-Cell Activation Antigen CD27,CD27 Molecule,CD27 Antigen,T Cell Activation Antigen S152,CD27L Receptor,TNFRSF7,S152,Tp55,T14.

### Description

Source : E.coli. CD27 Human Recombinant produced in E. coli is a single polypeptide chain containing 196 amino acids (21-191) and having a molecular mass of 21.8 kDa.CD27 is fused to a 25 amino acid His-tag at N-terminus & purified by proprietary chromatographic techniques. CD27 belongs to the TNF-receptor superfamily. CD27 is necessary for initiation and long-term maintenance of T cell immunity. CD27 binds to ligand CD70, and has a crucial role in regulating B-cell activation and immunoglobulin synthesis. The CD27 receptor transduces signals which result in the activation of NF-kappaB and MAPK8/JNK. Adaptor proteins TRAF2 and TRAF5 mediate the signaling process of CD27. CD27-binding protein (SIVA), which is a proapoptotic protein, can bind to the CD27 receptor and is believed to have a significant role in the apoptosis induced by CD27.

### Product Info

<b>Amount :</b>	20 µg
<b>Purification :</b>	Greater than 90% as determined by SDS-PAGE.
<b>Content :</b>	The CD27 solution (1mg/1ml) contains 20mM Tris-HCl buffer (pH 8.0), 0.4M Urea and 10% glycerol.
<b>Storage condition :</b>	Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Avoid multiple freeze-thaw cycles.
<b>Amino Acid :</b>	MGSSHHHHHH SSGLVPRGSH MGSHMTPAPK SCPERHYWAQ GKLCQMCEP GTFLVKDCDQ HRKAAQCDPC IPGVSFSPDH HTRPHCESCR HCNSGLLVRN CTITANAECA CRNGWQCRDK ECTECDPLPN PSLTARSSQA LSPHPQPTHL PYVSEMLEAR TAGHMQTLAD FRQLPARTLS THWPPQRS LC SSDFIR.

