

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-3835: Filamin Native Protein

Description

Source: Chicken Gizzard. Ultra Pure Filamin having a Molecular mass of 250 kDa. Filamin is a large (270kd) dimeric actin crosslinking protein from a variety of sources, which helps to stabilize the 3D cortical actin network. The fundamental structure of filamin is well conserved and consists of an actin binding domain at the N-terminus followed by a C-terminal rod domain consisting of numerous repeat segments ranging from 4 in C. elegans to 24 in mammalian cells. Each repeat in the rod domain consists of roughly 100 residues and forms an immunoglobulin like fold. Such immunoglobulin folds have been found in a variety of proteins and are responsible for protein-protein interactions. Filamin Human actin-binding protein (ABP), aka filamin, crosslinks actin filaments into orthogonal networks in cortical cytoplasm and participates in the anchoring of membrane proteins for the actin cytoskeleton. Mammalian filamin interacts directly with at least 30 proteins such as transmembrane receptors, second messenger-associated proteins, protein kinases, phosphatases and cytoskeletal proteins and these interactions have been shown to require one or more of the repeat elements in the rod domain.

Product Info

Amount: 50 µg

Purification : Greater than 90.0% as determined by SDS-PAGE.

Content:

The protein was lyophilized from a 1mg/ml solution containing 20mM Tris / acetate buffer pH 7.6,

0.1mM EDTA, 2mM DTT and 20mM NaCl. Each mg Filamin contains 550mg urea.

Lyophilized Filamin although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution Filamin should be stored at 4°C between 2-7 days and for future

use below -18°C.For long term storage it is recommended to add a carrier protein (0.1% HSA or

BSA).Please prevent freeze-thaw cycles.

Application Note

Storage condition:

It is recommended to reconstitute the lyophilized Filamin in sterile $18M\hat{l}$ ©-cm H2O not less than $100\hat{A}\mu g/ml$, which can then be further diluted to other aqueous solutions.

