

## 32-4005: ID2 Recombinant Protein

**Alternative Name** : DNA-binding protein inhibitor ID-2,bHLHb26,GIG8,ID2A,ID2H,MGC26389,Class B basic helix-loop-helix protein 26,Inhibitor of DNA binding 2.

### Description

Source : E.coli. ID2 Human Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 154 amino acids (1-134 a.a.) and having a molecular mass of 17kDa. ID2 is fused to a 20 amino acid His-tag at N-terminus & purified by proprietary chromatographic techniques. Inhibitor of DNA Binding 2 (ID2) is a part of the inhibitor of DNA binding family, whose members are transcriptional regulators that contain a helix-loop-helix (HLH) domain but not a basic domain. Members of the ID family inhibit the functions of basic helix-loop-helix transcription factors in a dominant-negative way by suppressing their heterodimerization partners through the HLH domains. ID2 play a role in negatively regulating cell differentiation and may be an inhibitor of tissue-specific gene expression.

### Product Info

**Amount :** 20 µg  
**Purification :** Greater than 85% as determined by SDS-PAGE.  
**Content :** ID2 protein solution (1mg/ml) contains 20mM Tris-HCl buffer (pH8.0), 10% glycerol and 0.4M Urea.  
**Storage condition :** Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Avoid multiple freeze-thaw cycles.  
**Amino Acid :** MGSSHHHHHH SSGLVPRGSH MKAFSPVRSV RKNLSLDHSL GISRSKTPVD DPMSLLYMN  
DCYSKLKELV PSIPQNKKVS KMEILQHVID YILDQLIALD SHPTIVSLHH QRPQONQASR TPLTTLNTDI  
SILSLQASEF PSELMNSDSK ALCG.