

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-4447: Recombinant Human Progesterone Receptor (Discontinued)

Alternative Name: Progesterone receptor, PR, Nuclear receptor subfamily 3 group C member 3, PGR, NR3C3.

Description

Source: Escherichia Coli. Progesterone Receptor Human Recombinant (a.a. 412-562) expressed in E.coli, shows a 43 kDa SDS-PAGE (including GST tag). The PGR is purified by proprietary chromatographic techniques. PGR belongs to the steroid receptor superfamily. PGR is located on chromosome 11q22. The progesterone receptor is an intracellular steroid receptor that specifically binds progesterone. The progesterone receptor (PgR) is an estrogen-regulated protein. PGR mediates the physiological effects of progesterone, which plays a fundamental role in reproductive events associated with establishing and maintaining pregnancy. PGR uses 2 distinct promotors and translation start sites in the first exon to produce 2 isoforms, A & B, which differ in their molecular weight (isoform A has 165 additional amino acids at the N-terminus). The progesterone receptor has an amino and a carboxyl terminal, the regulatory domain between them, a DNA binding domain, the hinge section, and the hormone binding domain. If no hormone is bound the carboxyl terminal inhibits transcription. Binding to a hormone stimulates a structural change which eliminates the inhibitory action. Following the progesterone binding to the receptor, restructuring with dimerization begins and the complex enters the nucleus where it binds to DNA.It has been proposed that expression of PgR determination indicates a responsive estrogen receptor (ER) pathway, and therefore, may predict likely response to endocrine therapy in human breast cancer. A number of studies have shown that PgR determination provides supplementary information to ER, both in predicting response to endocrine therapy and estimating survival. PgR has proved superior to ER as a prognostic indicator in some studies.

Product Info

Amount: 5 μ g

Content: PGR in 50mM Tris-HCl, pH7.5, 10mM L-glutathione (reduced).

Storage condition:

Storage condition:

Store vial at -20°C to -80°C. When stored at the recommended temperature, this protein is

stable for 12 months. Please prevent freeze-thaw cycles.

