

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-4729: Recombinant Human S100 Calcium Binding Protein A1

**Alternative Name** Protein S100-A1,S100 calcium-binding protein A1,S-100 protein alpha subunit,S-100 protein alpha chain,S100A1,S100A,S100,S100-alpha,S100-A1.

## **Description**

Source: Escherichia Coli. S100A1 Human Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 103 amino acids which include a 10 amino acid His Tag fused at N-terminus and having a total molecular mass of 11.66 kDa. S100A1 Human Recombinant is purified by proprietary chromatographic techniques. S100A1 is a member of the S100 family of calcium binding proteins with EF-hand type Ca+2 binding motive. S100A1 (Calcium Binding Protein A1) is involved in the activation of sarcoplasmatic calcium release and the regulation of intermediate filament polymerization. S100A1 may function in stimulation of Ca2+-induced Ca2+ release, inhibition of microtubule assembly, and inhibition of protein kinase C-mediated phosphorylation. Reduced expression of S100A1 has been implicated in cardiomyopathies.S100 proteins are localized either in the cytoplasm or the nucleus of a wide range of cells. There are at least 13 members in the S100 gene family, which are located as a cluster on chromosome 1q21.

## **Product Info**

Amount: 10 µg

**Purification:** Greater than 95.0% as determined by SDS-PAGE.

Content: The S100A1 protein was lyophilized from 0.4µm filtered solution at a concentration of 0.5mg/ml

containing 20mM Tris pH-7.5 and 50mM NaCl.

 $Lyophilized \ S100A1 \ although \ stable \ at \ room \ temperature \ for \ 3 \ weeks, \ should \ be \ stored \ desiccated$ 

Storage condition:

below -18°C. Upon reconstitution S100A1 should be stored at 4°C between 2-7 days and for future

use below -18°C.For long term storage it is recommended to add a carrier protein (0.1% HSA or

BSA). Please prevent freeze-thaw cycles.

Amino Acid: MKHHHHHHAS GSELETAMET LINVFHAHSG KEGDKYKLSK KELKELLQTE LSGFLDAQKD

VDAVDKVMKE LDENGDGEVD FQEYVVLVAA LTVACNNFFW ENS.

## **Application Note**

Add deionized water to a working concentration approximately 0.5mg/ml and let the lyophilized pellet dissolve completely. Product is not sterile! Please filter the product by appropriate sterile filter before using it in the cell culture.

