

## 32-4795: Recombinant Human E-Selectin HEK

**Alternative Name :** E-selectin, Endothelial leukocyte adhesion molecule 1, ELAM-1, Leukocyte-endothelial cell adhesion molecule 2, LECAM2, CD62E antigen, SELE, ELAM1, ELAM, ESEL, CD62E.

### Description

Source : HEK293 cells. SELE Human Recombinant produced by mammalian expression system in human cells is a single polypeptide chain containing 543 amino acids (22-556). SELE is fused to an 8 amino acid His-tag at C-terminus and is purified by proprietary chromatographic techniques. E-selectin which is also called Endothelial leukocyte adhesion molecule 1, ELAM1, ELAM belongs to a family of divalent cation-dependent carbohydrate-binding glycoproteins or adhesion molecules. Eselectin is expressed on the surface of endothelial cells and mediates the interaction of leukocytes and platelets with endothelial cells during an inflammatory response. E-selectin is present in single copy in the human genome and contains 14 exons spanning about 13 kb of DNA.

### Product Info

<b>Amount :</b>	10 µg
<b>Purification :</b>	Greater than 95% as determined by SDS-PAGE.
<b>Content :</b>	SELE was lyophilized from a 0.2 mM filtered solution of PBS and 4% Mannitol, pH 7.5.
<b>Storage condition :</b>	Lyophilized SELE although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution SELE should be stored at 4°C between 2-7 days and for future use below -18°C. Please prevent freeze-thaw cycles.
<b>Amino Acid :</b>	WSYNTSTEAMTYDEASAYCQQRHYTHLVAIQNKKEIEYLSILSYSPSYWIGIRKVNNVVWVGTKPLTEEA NWAPGEPNNRQKDEDCVEIYIKREKDVGMWNDERCSKKLALCYTAACNTNTSCSGHGECVETINNYTCKCDP GFGSLKCEQIVNCTALESPEHGLVCSHPLGNFSYNSSCSISCDRGYLPSSMETMQCMSSGEWSAPIACNVVE CDAVTNPANGFVECFQNPFSFPWNTTCTFDCEEGFELMGAQSLQCTSSGNWDNEKPTCKAVTCRAVRQPQN GSVRCSHSPAGEFTFKSSCNFTCEEGLQGPAAQVECTTQGGWTQIPVCEAFQCTALSNPERGYMNCPLSA SGSFRYGSCEFSCEQGFVLKGSRLQCGPTGEWDNEKPTCEAVRCDVHQPGLVRCASPIGEFTYKSSC AFSCEEGFELHGSTQLECTSQGQWTEEVPCQVVKCSSLAVPGKINMSCSGEPVFGTVCKFACPEGWTLNGS AARTCGATGHWSGLLPTCEAPTESNIPVDHHHHHH.

### Application Note

It is recommended to reconstitute the lyophilized SELE in 1xPBS to a concentration no less than 100 µg/ml, which can then be further diluted to other aqueous solutions.

