

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-4849: Recombinant Human Sirtuin-1

Alternative Name:

Sirtuin 1,SIRT1,SIR2L1,Regulatory Protein SIR2 Homolog 1,hSIR2,hSIRT1,SIR2-Like Protein

1,SIR2alpha,NAD-Dependent Deacetylase Sirtuin-1,NAD-Dependent Protein Deacetylase Sirtuin-1,Sir2-

Like 1, Sirtuin Type 1, EC 3.5.1.-.

Description

Source: Escherichia Coli. SIRT1 Human Recombinant produced in E.coli is a single, non-glycosylated polypeptide chain containing 280 amino acids (254-495) and having a molecular mass of 31.6 kDa.SIRT1 is fused to a 38 amino acid His-tag at Nterminus. Sirtuin-1 (SIRT1) belongs to the sirtuin family of proteins, homologs to the yeast Sir2 protein. Sirtuin family members are characterized by a sirtuin core domain and grouped into 4 classes. The functions of human sirtuins have not yet been established; though, yeast sirtuin proteins are known to regulate epigenetic gene silencing and suppress recombination of rDNA. It has been proposed that the human sirtuins may serve as intracellular regulatory proteins with mono-ADPribosyltransferase activity.

Product Info

Amount: 20 µg

Purification: "Greater than 90.0% as determined by SDS-PAGE."

The SIRT1 solution (0.5mg/ml) contains 20mM Tris-HCl buffer (pH 8.0), 10% glycerol and 0.4M Content:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of Storage condition:

time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please

avoid freeze thaw cycles.

MRGSHHHHHH GMASMTGGQQ MGRDLYDDDD KDRWGSHMKK IIVLTGAGVS VSCGIPDFRS Amino Acid:

> RDGIYARLAV DFPDLPDPQA MFDIEYFRKD PRPFFKFAKE IYPGQFQPSL CHKFIALSDK EGKLLRNYTQ NIDTLEQVAG IQRIIQCHGS FATASCLICK YKVDCEAVRG DIFNQVVPRC PRCPADEPLA IMKPEIVFFG ENLPEQFHRA MKYDKDEVDL LIVIGSSLKV RPVALIPSSI

PHEVPQILIN REPLPHLHFD VELLGDCDVI INELCHRLGG.

