

32-4852: Recombinant Human Sirtuin-5

Alternative Name :

Sirtuin 5, NAD-Dependent Lysine Demalonylase And Desuccinylase Sirtuin-5, Mitochondria, NAD-Dependent Protein Deacylase Sirtuin-5 Mitochondrial, Silent Mating Type Information Regulation 2 (S. Cerevisiae) Homolog 5, NAD-Dependent Deacetylase Sirtuin-5,

Description

Source : Escherichia Coli. SIRT5 Human Recombinant produced in E. Coli is a single, non-glycosylated polypeptide chain topological domain containing 300 amino acids (34-310 a.a) and having a molecular mass of 32.5 kDa. SIRT5 is fused to a 23 amino acid His-tag at N-terminus & purified by proprietary chromatographic techniques. SIRT5 belongs to the Sirtuin family of proteins, homologs to the yeast Sir2 protein. Sirtuin family members hold a sirtuin core domain and grouped into four classes. The protein encoded by this gene is included in class III of the sirtuin family. Human sirtuins functions were not yet determined; still, yeast sirtuin proteins are known to regulate epigenetic gene silencing and suppress recombination of rDNA. Studies propose that human sirtuin could operate as intracellular regulatory protein with mono-ADP-ribosyltransferase activity.

Product Info

Amount : 20 µg

Purification : "Greater than 90.0% as determined by SDS-PAGE."

Content : SIRT5 protein solution (0.5 mg/ml) containing 20 mM Tris-HCl buffer (pH 8.0), 0.15 M NaCl, 1 mM DTT and 30% glycerol.

Storage condition : Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please avoid freeze thaw cycles.

Amino Acid : MGSSHHHHHH SSGLVPRGSH MGSARPSSSM ADFRKFFAKA KHIVISGAG VSAESGVPTF
RGAGGYWRKW QAQDLATPLA FAHNPSRVWE FYHYRREVMG SKEPNAGHRA IAECETRLGK
QGRRVVVITQ NIDELHRKAG TKNLLEIHGS LFKTRCTSCG VVAENYKSPI CPALSGKGAP EPGTQDASIP
VEKLPRCEE A GCGLLRPHV VWFGENLDPA ILEEVDRELA HCDLCLVVG TSSVVYPAAMF APQVAARGVP
VAEFNTETTP ATNRFRFHFQ GPCGTTLPEA LACHENETVS

