

32-5032: Recombinant Human TRAF Family Member-Associated NFKB Activator

Alternative Name : TRAF,TRAF2,TRAF-interacting protein,ITRAF.

Description

Source : Escherichia Coli. TANK Human Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 448 amino acids (1-425a.a) and having a molecular mass of 50.2kDa. TANK is fused to a 23 amino acid His-tag at N-terminus & purified by proprietary chromatographic techniques. TRAF Family Member-Associated NFKB Activator (TANK) is located in the cytoplasm and binds Either TRAF1, TRAF2 or TRAF3. TANK is an inhibitor of TRAF function which regulates TRAF protein activity via sequestering TRAFs in a dormant position in the cytoplasm. Overexpression of TANK, inhibits TRAF2-mediated NF-Kappa-B activation signaled by CD40 and both TNF receptors and also inhibits LMP1-mediated NFKappa-B activation by blocking the connection of TRAF2 with LMP1.

Product Info

| | |
|----------------------------|--|
| Amount : | 20 µg |
| Purification : | Greater than 85.0% as determined by SDS-PAGE. |
| Content : | TANK protein solution (0.5mg/ml) contains 20mM Tris-HCl buffer (pH 8.0), 0.2M NaCl, 50% glycerol and 2mM DTT. |
| Storage condition : | A2LD1 should be stored desiccated below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles. |
| Amino Acid : | MGSSHHHHHH SSGLVPRGSH MGSMDKNIGE QLNKAYEAFR QACMDRDSAV KELQQKTENY EQRIREQEQ LSLQQTIDK LKSQLLLNS TQDNNGCVP LLEDSETRKN NLTLDQPQDK VISGIAREKL PKVRRQEVSS PRKETSARSL GSPLLHERGN IEKTFWDLKE EFHKICMLAK AQKDHLKLN IPDTATETQC SVPIQCTDKT DKQEALFKPQ AKDDINRGAP SITSVTPRGL CRDEEDTSFE SLSKFNKFP PMDNDSTFLH STPERGILS PATSEAVCQE KFNMEFRDNP GNFVKTEETL FEIQGIDPIA SAIQNLKTTD KTKPSNLVNT CIRTTLDRAA CLPPGDHNL YVNSFLLDP SDAPFSLDS PGKAIRGPQQ PIWKPFNPQD SDSVLSGTD SELHIPRVCE FCQAVFPPSI TSRGDFLRHL NSHFNGET. |