

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-5113: Recombinant Human Tubulin Polymerization Promoting Protein

Alternative Name Tubulin polymerization-promoting protein, TPPP, 25 kDa brain-specific protein, TPPP/p25, p24, p25-

alpha,TPPP1,p25,p25alpha.

## **Description**

Source: Escherichia Coli. TPPP Human Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 229 amino acids including a 10 a.a N-terminal His tag. The total molecular mass is 24.9kDa (calculated). Tubulin Polymerization Promoting Protein (TPPP) probably plays a part in the polymerization of tubulin into microtubules, as well as in microtubule bundling and the stabilization of existing microtubules, thus maintaining the integrity of the microtubule network. TPPP may also have a role in mitotic spindle assembly and nuclear envelope breakdown. TPPP/p235 level in cerebrospinal fluid is considerably higher in multiple sclerosis patients.

## **Product Info**

Amount: 10 µg

**Purification:** Greater than 85.0% as determined by SDS-PAGE.

Content: TPPP filtered (0.4µm) and lyophilized from 0.5mg/ml in 0.05M phosphate buffer and 0.075M NaCl,

pH 7.4.

Store lyophilized protein at -20°C. Aliquot the product after reconstitution to avoid repeated

Storage condition: freezing/thawing cycles. Reconstituted protein can be stored at 4°C for a limited period of time; it

does not show any change after two weeks at 4°C.

Amino Acid: MKHHHHHHASMADKAKPAKA ANRTPPKSPG DPSKDRAAKR LSLESEGAGE GAAASPELSA

LEEAFRRFAV HGDARATGRE MHGKNWSKLC KDCQVIDGRN VTVTDVDIVF SKIKGKSCRT ITFEQFQEAL EELAKKRFKD KSSEEAVREV HRLIEGKAPI ISGVTKAISS PTVSRLTDTT

KFTGSHKERF DPSGKGKGKA GRVDLVDESG YVSGYKHAGT YDQKVQGGK.

## **Application Note**

It is recommended to add 200µl deionized water to prepare a working stock solution of approximately 0.5mg/ml and let the lyophilized pellet dissolve completely. TPPP is not sterile! Please filter the product by an appropriate sterile filter before using it in the cell culture.

