

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-5164: Recombinant Human Thioredoxin-2

Alternative Name: Thioredoxin mitochondrial, Thioredoxin-2, TXN2, MTRX, TRX2, MT-TRX, TRX-2, TXN-2.

Description

Source: Escherichia Coli. MTRX Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 108 amino acids and having a molecular mass of 11 kDa. Thioredoxin-2 is a low molecular weight redox protein. TRX2 contains a redox active disulfide/dithiol group within the conserved Cys-Gly-Pro-Cys active site. The TXN2 is involved in the regulation of the mitochondrial membrane potential and in protection against oxidant-induced apoptosis. Upon stimulation of Fas, TXN2 mediates denitrosylation of mitochondria-associated caspase-3, a process required for caspase-3 activation, and promoted apoptosis. TRX2 is important at low oxidative stress conditions. MTRX is involved in the regulation of the mitochondrial membrane potential and cell death. Mitochondrial thioredoxin plays an important roles in protection against oxidant-induced apoptosis. Thioredoxin1 and thioredoxin2 have opposed regulatory functions on hypoxia-inducible factor-1alpha.

Product Info

Amount: 20 μg

Purification: Greater than 95.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.

Content: TXN2 protein solution contains 1x PBS pH-7.4.

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods

Storage condition: of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or

BSA). Avoid multiple freeze-thaw cycles.

Amino Acid: MTTFNIQDGP DFQDRVVNSE TPVVVDFHAQ WCGPCKILGP RLEKMVAKQH GKVVMAKVDI

DDHTDLAIEY EVSAVPTVLA MKNGDVVDKF VGIKDEDQLE AFLKKLIG.

Application Note

Specific activity is 3-4 A650/min/mg, obtained by measuring the increase of insulin precipitation in absorbance at 650 nm resulting from the reduction of insulin.

