

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 11-3045: Polyclonal Antibody to HMGB1

**Clonality:** Polyclonal

Application: WB
Reactivity: Human
Gene: HMGB1
Gene ID: 3146
Uniprot ID: P09429
Format: Purified

Alternative Name: HMGB1,HMG1 Isotype: Rabbit IgG

Immunogen Information: A partial length recombinant HMGB1 protein (amino acids 1-210) was used as the immunogen

for this antibody.

## **Description**

HMGB1 (High-mobility group protein B1) is a 25 kDa non-histone chromatin protein, secreted by immune cells through leaderless secretory pathway. HMGB1 supports transcription and also interacts with nucleosomes to loosen packed DNA and remodel the chromatin. The presence of HMGB1 in the nucleus depends on posttranslational modifications. When the protein is not acetylated, it stays in the nucleus, but hyperacetylation on lysine residues causes it to translocate into the cytosol. HMGB1 has the ability to bend DNA in order to facilitate binding of other proteins to the DNA. HMGB1 has one important role in V(D)J recombination where act as a cofactor of the RAG complex. HMGB1 stimulates a plethora of cellular responses like inflammation, attracting inflammatory cells, recruiting stem cells and promoting their proliferation. HMGB1 is highly expressed in human kidney, tonsils, adipose tissue, adrenal cortex and amniotic fluid.

## **Product Info**

**Amount :** 25 μg / 100 μg

**Purification:** Protein A Chromatography

Content: 25 µg in 50 µl/100 µg in 200 µl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium

azide is highly toxic.

**Storage condition :** Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid

repeated freeze and thaw cycles.

## **Application Note**

Western blot analysis: 1-2 µg/ml



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Fig-1: Western blot analysis of HMGB1. Anti-HMGB1 antibody (11-3045) was used at 1  $\mu\text{g/ml}$  on HepG2 lysate.