

36-1423: Monoclonal Antibody to MAP3K1 (Mitogen-Activated Protein Kinase Kinase Kinase 1)(Clone : 2F6)

Clonality : Monoclonal

Clone Name : 2F6

Application : WB,IHC

Reactivity : Human

Gene : MAP3K1

Gene ID : 4214

Uniprot ID : Q13233

Format : Purified

Alternative Name : MAP3K1,MAPKKK1,MEKK,MEKK1

Isotype : Mouse IgG2a, kappa

Immunogen Information : Partial recombinant MAP3K1 (aa1211-1310) (SKNSMTLDLNSSSKCDDSFSGCSSNSSNAVIPSDFTVFTP-VEEKRLDVNTELNSSIEDLLEASMPSSDITVTFKSEVAVLSPEKAENDDTYKDDVNHNQK)

Description

Mitogen-activated protein (MAP) kinase cascades are activated by various extracellular stimuli, including growth factors. The MEK kinases (also designated MAP kinase kinase kinases, MKKKs, MAP3Ks or MEKKs) phosphorylate and thereby activate the MEKs (also called MAP kinase kinases or MKKs), including ERK, JNK and p38. These activated MEKs in turn phosphorylate and activate the MAP kinases. The MEK kinases include Raf-1, Raf-B, Mos, MEK kinase-1, MEK kinase-2, MEK kinase-3, MEK kinase-4 and ASK 1 (MEK kinase- 5). MEK kinase-1 activates the ERK and c-Jun NH2-terminal kinase (JNK) pathways by phosphorylation of MAP2K1 and MAP2K4, and also activates the central protein kinases of the NFkB pathway, CHUK and IKKBK. Additionally, MEK kinase-1 uses an E3 ligase through its PHD domain, a RING-finger-like structure, to target proteins for degradation through ubiquitination.

Product Info

Amount : 100 µg


Purification : Affinity Chromatography

Content : 100 µg in 500 µl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly toxic.


Storage condition : Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid repeated freeze and thaw cycles.

Application Note


Western Blot (1-2ug/ml);Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris Buffer with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes)



Formalin-fixed, paraffin-embedded human Uterine Carcinoma stained with MAP3K1 Monoclonal Antibody (2F6).



Formalin-fixed, paraffin-embedded human Thyroid Carcinoma stained with MAP3K1 Monoclonal Antibody (2F6).



Formalin-fixed, paraffin-embedded human Cervical Carcinoma stained with MAP3K1 Monoclonal Antibody (2F6).