

## 31-1370: Goat Anti-Mouse Kappa-Biotin Conjugated

**Application :** IHC,IHC-P,IP,ELISA,FACS,WB,IHC-Fr

**Conjugate :** BIOTIN

**Isotype :** Goat IgG

### Product Info

**Amount :** 1.0 mg

**Purification :** Affinity chromatography on mouse kappa light chains covalently linked to agarose

**Content :** 0.5 mg/ml, 2.0 ml, Supplied in PBS/Sodium azide

**Storage condition :** Store at 2-8°C




Figure 1: ELISA plate was coated with purified mouse IgG<sup>1</sup>, IgM<sup>1</sup>, IgA<sup>1</sup>, IgG<sup>1</sup>, IgM<sup>1</sup>, and IgA<sup>1</sup>. Immunoglobulins were detected with serially diluted Goat Anti-Mouse Kappa-Biotin followed by Streptavidin-HRP .




Figure 2: Human pancreatic carcinoma cell line MIA PaCa-2 was stained with Mouse Anti-Cytokeratin 18 followed by Goat Anti-Mouse Kappa-Biotin, Streptavidin-FITC , and DAPI.




Figure 3: BALB/c mouse splenocytes were stained with Goat Anti-Mouse Kappa-Biotin followed by Streptavidin-FITC .




Figure 4: Human pancreatic carcinoma cell line MIA PaCa-2 was stained with Mouse Anti-Cytokeratin 18 followed by Goat Anti-Mouse Kappa-Biotin, Streptavidin-FITC and DAPI.