

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 10-6007: Monoclonal Antibody to IKK alpha (Clone: ABM10G9)

Clonality: Monoclonal **Clone Name:** ABM10G9 FACS.WB Application: Reactivity: Mouse, Human

Gene: **CHUK** Gene ID: 1147 **Uniprot ID:** 015111 Format: **Purified** 

Inhibitor of nuclear factor kappa-B kinase subunit alpha,I-kappa-B kinase alpha,Conserved

**Alternative Name:** helix-loop-helix ubiquitous kinase, Nuclear factor NF-kappa-B inhibitor kinase

alpha,NFKBIKA,Transcription factor 16

Isotype: Mouse IgG1, Kappa

Immunogen Information: Full length recombinant protein of IKK alpha was used as the immunogen for this antibody.

## **Product Info**

Amount: 25 μg / 100 μg

**Purification:** Protein G Chromatography

 $25~\mu g$  in  $50~\mu l/100~\mu g$  in  $200~\mu l$  PBS containing 0.05% BSA and 0.05% sodium azide. Sodium Content:

azide is highly toxic.

Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid Storage condition:

repeated freeze and thaw cycles.

## **Application Note**

WB: 2-4 μg/ml, FACS: 0.5-1 μg/10<sup>6</sup>




Figure-1: Western blot analysis of IKK alpha. Anti-IKK alpha antibody (Clone: ABM10G9) was used at 2 µg/ml on Hela, MCF7, U87, mSmall Intestine, PC3 and 293 lysates.





9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure-2: Intra cellular flow analysis of IKK alpha in Jurkat using  $0.5 \mu g/10^6$  cells of antibody (Clone: ABM10G9). Green represents isotype control; red represents anti-IKK alpha antibody. Goat anti-mouse PE conjugate was used as secondary antibody.




Figure-3: Intra cellular flow analysis of IKK alpha in 293 using 0.5  $\mu$ g/10^6 cells of antibody (Clone: ABM10G9). Green represents isotype control; red represents anti-IKK alpha antibody. Goat anti-mouse PE conjugate was used as secondary antibody.




Figure-4: Intra cellular flow analysis of IKK alpha in HeLa using 0.5  $\mu$ g/10^6 cells of antibody (Clone: ABM10G9). Green represents isotype control; red represents anti-IKK alpha antibody. Goat anti-mouse PE conjugate was used as secondary antibody.




Figure-5: Western blot analysis of IKK alpha. Anti-IKK alpha antibody (Clone: ABM10G9) was used at 2  $\mu$ g/ml on 3T3, Raw and EL-4 lysates.