

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

10-6007: Monoclonal Antibody to IKK alpha (Clone: ABM10G9)

Clone Name: ABM10G9
Application: FACS,WB
Reactivity: Mouse,Human

 Gene :
 CHUK

 Gene ID :
 1147

 Uniprot ID :
 015111

 Format :
 Purified

Inhibitor of nuclear factor kappa-B kinase subunit alpha,I-kappa-B kinase alpha,Conserved helix-

Alternative Name: loop-helix ubiquitous kinase, Nuclear factor NF-kappa-B inhibitor kinase

alpha,NFKBIKA,Transcription factor 16

Isotype: Mouse IgG1, Kappa

Immunogen Information: Full length recombinant protein of IKK alpha was used as the immunogen for this antibody.

Product Info

Amount : $25 \mu g / 100 \mu g$

Purification: Protein G Chromatography

Content: 25 μg in 50 μl/100 μg in 200 μl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium

azide is highly toxic.

Storage condition:

Storage condition:

Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid

repeated freeze and thaw cycles.

Application Note

WB: 2-4 µg/ml, FACS: 0.5-1 µg/10^6





Figure-1: Western blot analysis of IKK alpha. Anti-IKK alpha antibody (Clone: ABM10G9) was used at 2 μ g/ml on Hela, MCF7, U87, mSmall Intestine, PC3 and 293 lysates.








Figure-2: Intra cellular flow analysis of IKK alpha in Jurkat using 0.5 μ g/10^6 cells of antibody (Clone: ABM10G9). Green represents isotype control; red represents anti-IKK alpha antibody. Goat anti-mouse PE conjugate was used as secondary antibody.




Figure-3: Intra cellular flow analysis of IKK alpha in 293 using 0.5 μ g/10^6 cells of antibody (Clone: ABM10G9). Green represents isotype control; red represents anti-IKK alpha antibody. Goat anti-mouse PE conjugate was used as secondary antibody.




Figure-4: Intra cellular flow analysis of IKK alpha in HeLa using 0.5 μg/10^6 cells of antibody (Clone: ABM10G9). Green represents isotype control; red represents anti-IKK alpha antibody. Goat anti-mouse PE conjugate was used as secondary antibody.




Figure-5: Western blot analysis of IKK alpha. Anti-IKK alpha antibody (Clone: ABM10G9) was used at 2 μ g/ml on 3T3, Raw and EL-4 lysates.