

10-4088: Monoclonal Antibody to Mouse CD25 (Clone: PC61)

Clonality :	Monoclonal
Clone Name :	PC61
Application :	FACS
Reactivity :	Mouse
Gene :	ll2ra
Gene ID :	16184
Uniprot ID :	P01590
Format :	Purified
Alternative Name :	p55, ll2r
lsotype :	Rat IgG1
Immunogen Information : The B6.1 mouse cytotoxic T cell line was used as immunogen for this antibody.	

Description

CD25 is the alpha chain of the IL-2 receptor. It is a type I transmembrane protein present on activated T cells, activated B cells, some thymocytes, myeloid precursors, and oligodendrocytes that associates with CD122 to form a heterodimer that can act as a high-affinity receptor for IL-2. Though CD25 has been shown to influence effector function of lymphocytes, it is thought to play a greater role in immune tolerance in mice. Constitutive expression of CD25 on Treg cells is thought to be crucial to their survival and maintenance of immune homeostasis. Expression of CD25 and its role in immunology may be species dependent, since CD25 appears to play a larger role in T-cell effector responses in humans compared to mice, and may be somewhat dispensable for the maintenance of Treg cells as seen in patients treated with CD25-blocking antibodies.

Product Info

Amount : Purification :	25 μg / 100 μg Protein G Chromatography
Content :	25 μg in 50 μl/100 μg in 200 μl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly toxic.
Storage condition :	Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid repeated freeze and thaw cycles.

Application Note

FACS: 0.5-1 $\mu g/10\,{}^{\wedge}6$ cells

₩ abeomics

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982 Email: info@abeomics.com



Figure-1: Cell Surface flow staining of mCD25 on Con-A treated mouse splenocytes using 0.5 μ g/10^6 cells. Green represents isotype control; red represents anti-mouse CD25 antibody (10-4088, clone: PC61). Goat anti-rat FITC conjugated secondary antibody (ABEOMICS) was used.