

## 10-4154-NALE: NALE™ Monoclonal Antibody to hB7-2/CD86 (Clone: BU63) (No Azide Low Endotoxin)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	BU63
<b>Application :</b>	FACS
<b>Reactivity :</b>	Human
<b>Gene :</b>	CD86
<b>Gene ID :</b>	942
<b>Uniprot ID :</b>	P42081
<b>Format :</b>	Azide Free,Purified
<b>Alternative Name :</b>	Activation B7-2 antigen,B7-2 antigen, B-lymphocyte activation antigen B7-2, CD28 antigen ligand 2,CD28LG2B7-2 antigen, CTLA-4 counter-receptor B7.2, FUN-1, LAB72, MGC34413,T-lymphocyte activation antigen CD86
<b>Isotype :</b>	Mouse IgG1 Kappa
<b>Immunogen Information :</b>	ARH-77 (B-lymphoblastoid cell line) were used as immunogen for this antibody.

### Product Info

<b>Amount :</b>	25 µg / 100 µg
<b>Purification :</b>	Protein G Chromatography
<b>Content :</b>	25 µg in 50 µl/100 µg in 200 µl PBS containing 0.05% BSA. Azide free, low endotoxin.
<b>Storage condition :</b>	Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid repeated freeze and thaw cycles.

### Application Note

FACS: 1-2 µg/10<sup>6</sup> cells




Fig-1: Cell Surface flow analysis of hCD86 in PBMC (Granulocytes) using 1 µg/10<sup>6</sup> cells. Green represents isotype control (ABEOMICS); red represents anti-hCD86 antibody (10-4154). Goat anti-mouse PE conjugated secondary antibody (ABEOMICS) was used.




Fig-2: Cell Surface flow analysis of hCD86 in Raji using 1  $\mu\text{g}/10^6$  cells. Green represents isotype control (ABEOMICS); red represents anti-hCD86 antibody (10-4154). Goat anti-mouse PE conjugated secondary antibody (ABEOMICS) was used.




Fig-3: Cell Surface flow analysis of hCD86 in PBMC (Monocytes) using 0.5  $\mu\text{g}/10^6$  cells. Green represents isotype control (ABEOMICS); red represents anti-hCD86 antibody (10-4154). Goat anti-mouse PE conjugated secondary antibody (ABEOMICS) was used.