

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

35-1034: Polyclonal Antibody to Myc (Phospho-Thr358)

Clonality: Polyclonal Application: WB,IHC,IF Reactivity: Human Gene: MYC Gene ID: 4609 **Uniprot ID:** P01106 Format: Purified **Alternative Name:** c-myc Rabbit IgG Isotype:

Immunogen Information: Peptide sequence around phosphorylation site of threonine 358 (R-R-T(p)-H-N) derived from

Human Myc.

Description

Participates in the regulation of gene transcription. Binds DNA in a non-specific manner, yet also specifically recognizes the core sequence 5'-CAC[GA]TG-3'. Seems to activate the transcription of growth-related genes.

Product Info

Amount : 50 μl / 100 μl

Content: Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM

NaCl, 0.02% sodium azide and 50% glycerol.

Storage condition : Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid

repeated freeze and thaw cycles.

Application Note

Predicted MW: 60kd, Western blotting: $1:500\sim1:1000$, Immunohistochemistry: $1:50\sim1:100$, Immunofluorescence: $1:100\sim1:200$




Figure 1: Western blot analysis of extracts from HT29 cells untreated(lane 1) or treated with UV(lane 2) using Myc(Phospho-Thr358) Antibody 35-1034.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 2: Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using Myc(Phospho-Thr358) Antibody 35-1034 (left) or the same antibody preincubated with blocking peptide(right).




Figure 3: Immunofluorescence staining of methanol-fixed Hela cells using Myc(Phospho-Thr358) Antibody 35-1034.