

### 35-1048: Polyclonal Antibody to STAT5A (Phospho-Ser780)

<b>Clonality :</b>	Polyclonal
<b>Application :</b>	WB,IHC
<b>Reactivity :</b>	Human,Mouse,Rat
<b>Gene :</b>	STAT5A
<b>Gene ID :</b>	6776
<b>Uniprot ID :</b>	P42229
<b>Format :</b>	Purified
<b>Alternative Name :</b>	MGF, MPF, Mammary gland factor, STA5A,STAT5
<b>Isotype :</b>	Rabbit IgG
<b>Immunogen Information :</b>	Peptide sequence around phosphorylation site of serine 780 (R-L-S(p)-P-P) derived from Human STAT5A.

#### Description

Carries out a dual function: signal transduction and activation of transcription. Binds to the GAS element and activates PRL-induced transcription.

#### Product Info

<b>Amount :</b>	50 µl / 100 µl
<b>Content :</b>	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg <sup>2+</sup> and Ca <sup>2+</sup> ), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
<b>Storage condition :</b>	Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid repeated freeze and thaw cycles.

#### Application Note

Predicted MW: 90kd, Western blotting: 1:500~1:1000, Immunohistochemistry: 1:50~1:100




Figure 1: Western blot analysis of extracts from K562 cells untreated or treated with IL-4 using STAT5A(Phospho-Ser780) Antibody 35-1048 .




Figure 2: Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using STAT5A(Phospho-Ser780) Antibody 35-1048 (left) or the same antibody preincubated with blocking peptide(right).