

35-1148: Polyclonal Antibody to PAK1 (Phospho-Thr212)

| | |
|--------------------------------|-------------------------------------------------------------------------------------------------------|
| Clonality : | Polyclonal |
| Application : | WB,IHC,IF |
| Reactivity : | Human,Mouse,Rat |
| Gene : | PAK1 |
| Gene ID : | 5058 |
| Uniprot ID : | Q13153 |
| Format : | Purified |
| Alternative Name : | p21-activated kinase 1, PAK-1, p65-PAK, Alpha-PAK |
| Isotype : | Rabbit IgG |
| Immunogen Information : | Peptide sequence around phosphorylation site of threonine 212 (P-V-T(p)-P-T) derived from Human PAK1. |

Description

The activated kinase acts on a variety of targets. Likely to be the GTPase effector that links the Rho-related GTPases to the JNK MAP kinase pathway. Activated by CDC42 and RAC1. Involved in dissolution of stress fibers and reorganization of focal complexes. Involved in regulation of microtubule biogenesis through phosphorylation of TBCB. Activity is inhibited in cells undergoing apoptosis, potentially due to binding of CDC2L1 and CDC2L2.

Product Info

| | |
|----------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Amount : | 50 µl / 100 µl |
| Content : | Supplied at 1.0mg/mL in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. |
| Storage condition : | Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid repeated freeze and thaw cycles. |

Application Note

Predicted MW: 68kd, Western blotting: 1:500~1:1000, Immunohistochemistry: 1:50~1:100, Immunofluorescence: 1:100~1:200




Figure 1: Western blot analysis of extracts from 293 cells untreated or treated with forskolin using PAK1(Phospho-Thr212) Antibody 35-1148 .




Figure 2: Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using PAK1(Phospho-Thr212) Antibody 35-1148 (left) or the same antibody preincubated with blocking peptide(right).




Figure 3: Immunofluorescence staining of methanol-fixed HeLa cells using PAK1(Phospho-Thr212) Antibody 35-1148 .