

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 35-1153: Polyclonal Antibody to Zap-70 (Phospho-Tyr493)

Clonality: Polyclonal Application: WB,IHC

**Reactivity:** Human, Mouse, Rat

Gene: ZAP70
Gene ID: 7535
Uniprot ID: P43403
Format: Purified

Alternative Name: 70 kDa zeta-associated protein, SRK, Syk-related tyrosine kinase, ZA70, ZAP-70

**Isotype:** Rabbit IgG

Immunogen Information: Peptide sequence around phosphorylation site of tyrosine 493 (S-Y-Y(p)-T-A) derived from

Human Zap-70.

## **Description**

Plays a role in T-cell development and lymphocyte activation. Essential for TCR-mediated IL-2 production. Isoform 1 induces TCR-mediated signal transduction, isoform 2 does not.

## **Product Info**

**Amount :** 50 μl / 100 μl

Content: Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM

NaCl, 0.02% sodium azide and 50% glycerol.

**Storage condition :** Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid

repeated freeze and thaw cycles.

## **Application Note**

Predicted MW: 70kd, Western blotting: 1:500~1:1000, Immunohistochemistry: 1:50~1:100




Figure 1: Western blot analysis of extracts from JK cells untreated or treated with anisomycin using Zap-70(Phospho-Tyr493) Antibody 35-1153.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 2: Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using Zap-70(Phospho-Tyr493) Antibody 35-1153 (left) or the same antibody preincubated with blocking peptide(right).