

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 35-1155: Polyclonal Antibody to IkB- Alpha (Phospho-Tyr42) Antibody

Clonality: Polyclonal
Application: IHC,WB
Reactivity: Mouse,Human

Gene : NFKBIA
Gene ID : 4792
Uniprot ID : P25963
Format : Purified

Alternative Name: IKBA, MAD-3, NFKBI, NFKBIA

**Isotype:** Rabbit IgG

Immunogen Information: Peptide sequence around phosphorylation site of tyrosine 42 (E-E-Y(p)-E-Q) derived from

Human IkB-a

## **Description**

Inhibits the activity of dimeric NF-kappa-B/REL complexes by trapping REL dimers in the cytoplasm through masking of their nuclear localization signals. On cellular stimulation by immune and proinflammatory responses, becomes phosphorylated promoting ubiquitination and degradation, enabling the dimeric RELA to transcript to the nucleus and activate transcription.

## **Product Info**

**Amount :** 50 μl / 100 μl

Content: Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM

NaCl, 0.02% sodium azide and 50% glycerol.

**Storage condition :** Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid

repeated freeze and thaw cycles.

## **Application Note**

Predicted MW: 39kd, Western blotting: 1:500~1:1000, Immunohistochemistry: 1:50~1:100




Figure 1: Western blot analysis of extract from Hela cells untreated or treated with serum using IkB-a(Phospho-Tyr42) Antibody 35-1155 .



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 2: Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using IkB-a(phospho-Tyr42) antibody(35-1155).