

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

35-1158: Polyclonal Antibody to PAK1/PAK2/PAK3 (Phospho-Thr423/Thr402/Thr421)

Clonality: Polyclonal Application: IHC,IF

Reactivity: Rat, Mouse, Human

Gene: PAK1 Gene ID: 5058 **Uniprot ID:** Q13153 Format: Purified

Alternative Name: p21-activated kinase 1, PAK-1, p65-PAK, Alpha-PAK

Isotype: Rabbit IgG

Peptide sequence around phosphorylation site of threonine 423 /402/421 (R-S-T(p)-M-V) derived Immunogen Information:

from Human PAK1/PAK2/PAK3.

Description

The activated kinase acts on a variety of targets. Likely to be the GTPase effector that links the Rho-related GTPases to the JNK MAP kinase pathway. Activated by CDC42 and RAC1. Involved in dissolution of stress fibers and reorganization of focal complexes. Involved in regulation of microtubule biogenesis through phosphorylation of TBCB. Activity is inhibited in cells undergoing apoptosis, potentially due to binding of CDC2L1 and CDC2L2. Parsons M, et al. (2005) Mol Cell Biol; 25:1680-1695. Jung JH, et al. (2005) J Biol Chem; 280:40025-40031. Lampugnani MG, et al. (2002) Mol Biol Cell; 13: 1175-1189 Wang RA, et al. (2002) EMBO J; 21: 5437 - 5447.

Product Info

Amount: $50 \mu l / 100 \mu l$

Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM Content:

NaCl, 0.02% sodium azide and 50% glycerol.

Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid Storage condition:

repeated freeze and thaw cycles.

Application Note

Predicted MW: 61-67 (PAK2) 68-74 (PAK1/3)kd, Immunohistochemistry: 1:50~1:100, Immunofluorescence: 1:100~1:200




Figure 1: Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using PAK1/PAK2/PAK3(Phospho-Thr423/Thr402/Thr421) Antibody 35-1158 (left) or the same antibody preincubated with blocking peptide(right).



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 2: Immunofluorescence staining of methanol-fixed Hela cells using PAK1/PAK2/PAK3(Phospho-Thr423/Thr402/Thr421) Antibody 35-1158 .